K. T. S. P. Mandal's Hutatma Rajguru Mahavidyalaya Rajgurunagar 410505

Department of zoology

T. Y. B. Sc

ZO 353: Biological chemistry Semester V


> By Prof. P. P. Shindekar

> > 1

Introduction to Proteinsand Amino Acids


Introduction

- Twenty percent of the human body is made up of proteins. Proteins are the large, complex molecules that are critical fornormal functioning of cells.
- They are essential for the structure, function, and regulation of the body's tissues and organs.
- Proteins are made up of smaller units called amino acids, which are building blocks of proteins. They are attached to one anotherby peptide bonds forming a long chain of proteins.




Amino acid structure and its classification

- An amino acid contains both a carboxylic group and an amino group. Amino acids that have an amino group bonded directly tothe alpha-carbon are referred to as **alpha amino acids**.
- Every alpha amino acid has a carbon atom, called an alpha carbon, Cα; bonded to a carboxylic acid, –COOH group; anamino, –NH2 group; a hydrogen atom; and an R group that isunique for every amino acid.



Classification of amino acids

• There are 20 amino acids. Based on the nature of their 'R' group, they are classified based on their polarity as:



Classification based on essentiality:

Essential amino acids are the amino acids which you need through your diet because your body cannot make them. Whereas **non essential aminoacids** are the amino acids which are not an essential part of your diet because they can be synthesized by your body.

Essential


Histidine Isoleucine Leucine Methionine Phenyl alanine Threonine Tryptophan Valine

Non essential

Alanine Arginine Aspargine Aspartate Cystine Glutamic acid Glycine Ornithine Proline Serine Tyrosine

Peptide bonds

- Amino acids are linked together by 'amide groups' calledpeptide bonds.
- During protein synthesis, the carboxyl group of amino acidat the end of the growing polypeptide chain reacts with theamino group of an incoming amino acid, releasing a molecule of water. The resulting bond between the amino acids is a peptide bond.




Structure of proteins

- The sequence of a protein is determined by the DNA of the gene that encodes the protein (or thatencodes a portion of the protein, for multi- subunit proteins).
- A change in the gene's DNA sequence may lead to a change in the amino acid sequence of the protein. Even changing just one amino acid in a protein's sequence can affect the protein's overallstructure and function.
- To understand how a protein gets its final shape or conformation, we need to understand the fourlevels of protein structure: primary, secondary, tertiary, and quaternary


Primary Structure

- The simplest level of protein structure, primary structure is simply the sequence of amino acids in apolypeptide chain.
- The hormone insulin has two polypeptide chains A, and B. The sequence of the A chain, and the sequence of the B chain can be considered as an example for primary structure.




Secondary structure

- **secondary structure**, refers to local folded structures that formwithin a polypeptide due to interactions between atoms.
- The most common types of secondary structures are the α helix and the β pleated sheet. Both structures are held in shape by hydrogen bonds, which form between the carbonyl O of one aminoacid and the amino H of another.




Tertiary structure

- The overall three-dimensional structure of a polypeptide is called its**tertiary structure**. The tertiary structure is primarily due to interactions between the R groups of the amino acids that make upthe protein.
- Important to tertiary structure are **hydrophobic interactions**, in which amino acids with nonpolar, hydrophobic R groups cluster together on the inside of the protein, leaving hydrophilic amino acidson the outside to interact with surrounding water molecules.
- Also, **Disulfide bonds**, covalent linkages between the sulfur- containing side chains of cysteines, are much stronger than the othertypes of bonds that contribute to tertiary structure



Quaternary structure

- When multiple polypeptide chain subunits come together, then the protein attains its quaternarystructure.
- An example for quaternary structure is hemoglobin. The hemoglobin carries oxygen in the blood and is made up of four subunits, two each of the α and β types.



Denaturation and protein folding

- Each protein has its own unique shape. If the temperature or pH of a protein's environment ischanged, or if it is exposed to chemicals, these interactions may be disrupted, causing the protein to lose its three-dimensional structure and turn back into an unstructured string of amino acids.
- When a protein loses its higher-order structure, but not its primary sequence, it is said to be **denatured**. Denatured proteins are usually non-functional.

